arXiv:math/0703476v1 [math.NT] 16 Mar 2007

A NOTE ON THE q-GENOCCHI NUMBERS AND POLYNOMIALS

Taekyun Kim

Abstract. In this paper we discuss the new concept of the q-extension of Genocchi numbers and give the some relations between q-Genocchi polynomials and q-Euler numbers.

§1. Introduction

A number given by the generating function

$$\frac{2t}{e^t+1}=\sum_{n=0}^{\infty}G_n\frac{t^n}{n!}. \tag{1}$$

It satisfies $G_1=1, G_3=G_4=G_7=\cdots=0$, and even coefficients are given by $G_{2m}=2(1-2^{2n})B_{2n}=2nE_{2n-1}(0)$, where B_n is a Bernoulli number and $E_n(x)$ is an Euler polynomial. The first few Genocchi numbers for $n=2,4,\cdots$ are $-1,-3,17,-155,2073,\cdots$. The Euler polynomials are well known as

$$\frac{2}{e^t+1}e^{xt}=\sum_{n=0}^{\infty}E_n(x)\frac{t^n}{n!},\ \text{see [1, 3, 7, 8, 9]}. \tag{2}$$

By (1) and (2) we easily see that

$$E_n(x)=\sum_{k=0}^{n}\binom{n}{k}\frac{G_{k+1}}{k+1}x^{n-k},\ \text{where}\ \binom{n}{k}=\frac{n(n-1)\cdots(n-k+1)}{k!},\ \text{cf. [4, 5, 6]}. \tag{3}$$

For $m,n\geq 1$ and m odd, we have

$$(n^m-n)G_m=\sum_{k=1}^{m-1}\binom{m}{k}n^kG_kZ_{m-k}(n-1), \tag{4}$$

Key words and phrases. p-adic q-integrals, Genocchi numbers, q-Genocchi numbers, polynomials, sums of powers.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$-TEX

where $Z_m(n) = 1^m - 2^m + 3^m - \cdots + (-1)^{n-1}n^m$, see[3]. From (10 we derive

$$2t = \sum_{n=0}^{\infty} \left((G+1)^n + G_n\right) \frac{t^n}{n!},$$

where we use the technique method notation by replacing G^m by $G_m (m \geq 0)$, symbolically. By (5), we see that

$$G_0 = 0, \quad (G+1)^n - G_n = \begin{cases} 2 & \text{if } n = 1 \\ 0 & \text{if } \ n > 1. \end{cases} \tag{6}$$

Let p be a fixed odd prime, and let $\mathbb{C}_p$ denote the p-adic completion of the algebraic closure of $\mathbb{Q}_p$(=p-adic number field). For d a fixed positive integer with $(p, d) = 1$, let

$$\begin{aligned} X = X_d &= \varprojlim_{N} \mathbb{Z}/dp^N\mathbb{Z}, \\ X_1 &= \mathbb{Z}_p, \\ X^* &= \bigcup_{\substack{0<a<dp \\ (a,p)=1}} (a + dp\mathbb{Z}_p), \\ a + dp^N\mathbb{Z}_p &= \{x \in X | x \equiv a \pmod{dp^n}\}, \end{aligned}$$

where $a \in \mathbb{Z}$ lies in $0 \leq a < dp^N$.

Ordinary q-calculus is now very well understood from many different point of views. Let us consider a complex number $q \in \mathbb{C}$ with $|q| < 1$ (or $q \in \mathbb{C}_p$ with $|1-q|_p < p^{-\frac{1}{p-1}}$) as an indeterminate. The q-basic numbers are defined by

$$[x]_q = \frac{q^x - 1}{q - 1} = 1 + q + q^2 + \cdots + q^{x-1},$$

and

$$[x]_{-q} = \frac{-(-q)^x + 1}{q + 1} = 1 - q + q^2 + \cdots + (-q)^{x-1}.$$

We say that f is a uniformly differentiable function at a point $a \in \mathbb{Z}_p$ and denote this property by $f \in UD(\mathbb{Z}_p)$, if the difference quotients

$$F_f(x, y) = \frac{f(x) - f(y)}{x - y}$$

have a limit $l = f'(a)$ as $(x, y) \to (a, a)$.

For $f \in UD(\mathbb{Z}_p)$, let us start with the expression

$$\frac{1}{[p^N]_q} \sum_{0 \leq j < p^N} q^j f(j) = \sum_{0 \leq j < p^N} f(j)\mu_q(j + p^N\mathbb{Z}_p),$$

representing a q-analogue of Riemann sums for f, cf. [5]. The integral of f on $\mathbb{Z}_p$ will be defined as limit ($n \to \infty$) of those sums, when it exists. The p-adic q-integral of the function $f \in UD(\mathbb{Z}_p)$ is defined by

$$I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N\to\infty} \frac{1}{[p^N]_q} \sum_{0\le x<p^N} f(x) q^x, \quad \text{(see [5, 10, 11, 12]).}$$

In the previous paper [4] author constructed the q-extension of Euler polynomials in the fermionic sense of p-adic q-integral at $q = -1$ as follows:

$$(7) \quad E_{n,q}(x) = \int_{\mathbb{Z}_p} [t+x]_q^n d\mu_{-q}(t), \text{ where } \mu_{-q}(x + p^N\mathbb{Z}_p) = \lim_{q\to -q} \mu_q(x + p^N\mathbb{Z}_p).$$

From (7), we note that

$$(8) \qquad E_{n,q}(x) = \frac{[2]_q}{(1-q)^n} \sum_{l=0}^{n} \binom{n}{l} \frac{(-1)^l}{1+q^{l+1}} q^{lx}, \text{ see [4].}$$

The q-extension of Genocchi numbers is defined as

$$(9) \qquad g_q^*(t) = [2]_q t \sum_{n=0}^{\infty} (-1)^n q^n e^{[n]_q t} = \sum_{n=0}^{\infty} G_{n,q}^* \frac{t^n}{n!}, \text{ see [4].}$$

The following formula is well known in [4, 7]:

$$(10) \qquad E_{n,q}(x) = \sum_{k=0}^{n} \binom{n}{k} [x]_q^{n-k} q^{kx} \frac{G_{k+1,q}^*}{k+1}.$$

The modified q-Euler numbers are defined as

$$\xi_{0,q} = \frac{[2]_q}{2}, \quad (q\xi+1)^k + \xi_{k,q} = \begin{cases} [2]_q & \text{if } k = 0 \\ 0 & \text{if } \ k \neq 0, \end{cases}$$

with the usual convention of replacing ξ^i by $\xi_{i,q}$, see [10]. Thus, we derive the generating function of $\xi_{n,q}$ as follows:

$$(11) \qquad F_q(t) = [2]_q \sum_{k=0}^{\infty} (-1)^k e^{[k]_q t} = \sum_{n=0}^{\infty} \xi_{n,q} \frac{t^n}{n!}.$$

Now we also consider the q-Euler polynomials $\xi_{n,q}(x)$ as

$$(12) \qquad F_q(t,x) = [2]_q \sum_{k=0}^{\infty} (-1)^k e^{[k+x]_q t} = \sum_{n=0}^{\infty} \xi_{n,q}(x) \frac{t^n}{n!}.$$

From (12) we note that

$$\xi_{n,q}(x) = \sum_{l=0}^{n} \binom{n}{l} \xi_{l,q} q^{lx} [x]_q^{n-l}, \text{ see [10]}. \tag{13}$$

In the recent several authors studied the q-extension of Genocchi numbers and polynomials (see[1, 2, 5, 6, 7, 12]). In this paper we discuss the new concept of the q-extension of Genocchi numbers and give the some relations between q-Genocchi numbers and q-Euler numbers.

§2. q-extension of Genocchi numbers

In this section we assume that $q \in \mathbb{C}$ with $|q| < 1$. Now we consider the q-extension of Genocchi numbers as follows:

$$g_q(t) = [2]_q t \sum_{k=0}^{\infty} (-1)^k e^{[k]_q t} = \sum_{n=0}^{\infty} G_{n,q} \frac{t^n}{n!}. \tag{14}$$

In Eq.(14), it is easy to show that $\lim_{q \to 1} g_q(t) = \frac{2t}{e^t+1} = \sum_{n=0}^{\infty} G_n \frac{t^n}{n!}$. From (14) we derive

$$\begin{aligned} g_q(t) &= [2]_q t \sum_{k=0}^{\infty} (-1)^k \sum_{m=0}^{\infty} [k]_q^m \frac{t^m}{m!} = [2]_q \sum_{k=0}^{\infty} (-1)6k \sum_{m=1}^{\infty} m [k]_q^{m-1} \frac{t^m}{m!} \\ &= [2]_q \sum_{k=0}^{\infty} (-1)^k \sum_{m=0}^{\infty} m [k]_q^{m-1} \frac{t^m}{m!}. \end{aligned} \tag{15}$$

By (15), we easily see that

$$g_q(t) = [2]_q \sum_{m=0}^{\infty} \left(m \left(\frac{1}{1-q} \right)^{m-1} \sum_{l=0}^{m-1} \binom{m-1}{l} (-1)^l \frac{1}{1+q^l} \right) \frac{t^m}{m!}. \tag{16}$$

From (14) and (16) we note that

$$\sum_{m=0}^{\infty} G_{m,q} \frac{t^m}{m!} = \sum_{m=0}^{\infty} \left(m [2]_q \left(\frac{1}{1-q} \right)^{m-1} \sum_{l=0}^{m-1} \binom{m-1}{l} \frac{(-1)^l}{1+q^l} \right) \frac{t^m}{m!}. \tag{17}$$

By comparing the coefficients on both sides in Eq.(17), we have the following theorem:

Theorem 1. *For $m \geq 0$ we have*

$$G_{m,q} = m[2]_q \left(\frac{1}{1-q} \right)^{m-1} \sum_{l=0}^{m-1} \binom{m-1}{l} \frac{(-1)^l}{1+q^l}.$$

From Theorem 1, we easily derive the following corollary:

Corollary 2. *For $n \in \mathbb{N}$ we have*

$$G_{0,q} = 0, \quad (qG+1)^k + G_{k,q} = \begin{cases} \frac{[2]_q^2}{2} & \text{if } k = 1, \\ 0 & \text{if } \ k > 1, \end{cases}$$

with the usual convention of replacing G^i by $G_{i,q}$.

Remark. We note that Corollary 2 is the q-extension of Eq.(6). By (10) and Corollary 2, we obtain the following theorem:

Theorem 3. *For $n \in \mathbb{N}$ we have*

$$\xi_{n,q} = \frac{G_{n+1,q}}{n+1}.$$

From (12) we derive

$$\begin{aligned} F_q(x,t) &= [2]_q \sum_{n=0}^{\infty} (-1)^n e^{[n+x]_q t} = q^x t \frac{[2]_q}{q^x t} e^{[x]_q t} \sum_{n=0}^{\infty} (-1)^n e^{q^x [n]_q t} \\ &= e^{[x]_q t} \sum_{n=0}^{\infty} q^{nx} \frac{G_{n+1,q}}{n+1} \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left(\sum_{k=0}^{n} \binom{n}{k} [x]_q^{n-k} q^{kx} \frac{G_{k+1,q}}{k+1} \right) \frac{t^n}{n!}. \quad (18) \end{aligned}$$

By (18), we easily see that

$$\xi_{n,q}(x) = \sum_{k=0}^{n} \binom{n}{k} [x]_q^{n-k} q^{kx} \frac{G_{k+1,q}}{k+1}.$$

This formula can be considered as the q-extension of Eq.(3). Let us consider the q-analogue of Genocchi polynomials as follows:

$$(19) \qquad g_q(x,t) = [2]_q t \sum_{k=0}^{\infty} (-1)^k e^{[k+x]_q t} = \sum_{n=0}^{\infty} G_{n,q}(x) \frac{t^n}{n!}.$$

Thus, we note that $\lim_{q\to 1} g_q(x,t) = \frac{2t}{e^t+1}e^{xt} = \sum_{n=0}^{\infty} G_n(x)\frac{t^n}{n!}$. From Eq.(19), we easily derive

$$G_{n,q}(x) = [2]_q n \left(\frac{1}{1-q}\right)^{n-1} \sum_{l=0}^{n-1} \frac{(-1)^l}{1+q^l} q^{lx} \binom{n-1}{l}. \tag{20}$$

By (19) we also see that

$$\begin{aligned}
\sum_{n=0}^{\infty} G_{n,q}(x)\frac{t^n}{n!} &= [2]_q t \sum_{k=0}^{\infty} (-1)^k e^{[k+x]_q t} = [2]_q t \sum_{a=0}^{m-1} (-1)^a \sum_{k=0}^{\infty} (-1)^k e^{[k+\frac{a+x}{m}]_{q^m}[m]_q t} \\
&= \frac{[2]_q}{[m]_q [2]_{q^m}} \sum_{a=0}^{m-1} (-1)^a \left([m]_q t [2]_{q^m} \sum_{k=0}^{\infty} (-1)^k e^{[m]_q t [k+\frac{a+x}{m}]_{q^m}} \right) \\
&= \sum_{n=0}^{\infty} \left(\frac{[2]_q}{[m]_q [2]_{q^m}} \sum_{a=0}^{m-1} (-1)^a [m]_q^n G_{n,q^m}\left(\frac{x+a}{m}\right) \right) \frac{t^n}{n!} \\
&= \sum_{n=0}^{\infty} \left(\frac{[2]_q}{[2]_{q^m}} [m]_q^{n-1} \sum_{a=0}^{m-1} (-1)^a G_{n,q^m}\left(\frac{x+a}{m}\right) \right) \frac{t^n}{n!}, \text{ where } m \in \mathbb{N} \text{ odd.}
\end{aligned}$$

Therefore we obtain the following theorem:

Theorem 4. *Let $m(=\mathit{odd}) \in \mathbb{N}$. Then we have the distribution of the q-Genocchi polynomials as follows:*

$$G_{n,q}(x) = \frac{[2]_q}{[2]_{q^m}} [m]_q^{n-1} \sum_{a=0}^{m-1} (-1)^a G_{n,q^m}\left(\frac{x+a}{m}\right),$$

where n is positive integer.

Theorem 4 will be used to construct the p-adic q-Genocchi measures which will be treated in the next section. Let χ be a primitive Dirichlet character with a conductor $d(=\mathit{odd}) \in \mathbb{N}$. Then the generalized q-Genocchi numbers attached to χ are defined as

$$g_{\chi,q}(t) = [2]_q t \sum_{a=0}^{d-1} \chi(n)(-1)^n e^{[n]_q t} = \sum_{n=0}^{\infty} G_{n,\chi,q} \frac{t^n}{n!}. \tag{21}$$

From (21) we derive

$$G_{n,\chi,q} = \frac{[2]_q}{[2]_{q^d}} [d]_q^{n-1} \sum_{a=0}^{d-1} (-1)^a \chi(a) G_{n,q^d}(\frac{a}{d}). \tag{22}$$

§3. p-adic q-Genocchi measures

In this section we assume that $q \in \mathbb{C}_p$ with $|1-q|_p < p^{-\frac{1}{p-1}}$ so that $q^x = \exp(x \log q)$. Let χ be a primitive Dirichlet's character with a conductor $d(= odd) \in \mathbb{N}$. For any positive integers N, k and $d(= odd)$, let $\mu_k = \mu_{k,q;G}$ be defined as

$$\mu_k(a + dp^N\mathbb{Z}_p) = (-1)^a [dp^N]_q^{k-1} \frac{[2]_q}{[2]_{q^{dp^N}}} G_{k,q^{dp^N}}(\frac{a}{dp^N}). \tag{23}$$

By using Theorem 4 and (23), we show that

$$\sum_{i=0}^{p-1} \mu_k(a + idp^N + dp^{N+1}\mathbb{Z}_p) = \mu_k(a + dp^N\mathbb{Z}_p).$$

Therefore we obtain the following theorem:

Theorem 5. *Let d be odd positive integer. For any positive integers N, k, and let $\mu_k = \mu_{k,q;G}$ be defined as*

$$\mu_k(a + dp^N\mathbb{Z}_p) = (-1)^a [dp^N]_q^{k-1} \frac{[2]_q}{[2]_{q^{dp^N}}} G_{k,q^{dp^N}}(\frac{a}{dp^N}).$$

Then μ_k can be extended to a distribution on X.

From the definition of μ_k and (22) we note that

$$\int_X \chi(x) d\mu_k(x) = G_{k,\chi,q}.$$

By (14) and (16), it is not difficult to show that

$$G_{n,q}(x) = \sum_{k=0}^{n} \binom{n}{k} [x]_q^{n-k} q^{kx} G_{k,q}. \tag{24}$$

From (23) and (24) we derive

$$d\mu_k(a) = \lim_{N\to\infty} \mu_k(a + dp^N\mathbb{Z}_p) = k[a]_q^{k-1} d\mu_{-q}(a). \tag{25}$$

Therefore we obtain the following corollary:

Corollary 6. *Let k be a posotove integer. Then we have*

$$G_{k,\chi,q} = \int_X \chi(x) d\mu_k(x) = k \int_X \chi(x)[x]_q^{k-1} d\mu_{-q}(x).$$

Moreover,

$$G_{k,q} = k \int_X [x]_q^{k-1} d\mu_{-q}(x).$$

Remark. In the recent paper(see [1]) Cenkci-Can-Kurt have studied q-Genocchi numbers and polynomials and p-adic q-Genocchi measures. Our q-Genocchi numbers and polynomials to treat in this paper are different their q-Genocchi numbers and polynomials.

REFERENCES

[1] M. Cenkci, M. Can and V. Kurt, *q-extension of Genocchi Numbers*, J. Korean Math. Soc. **43** (2006), 183–198.

[2] M. Cenkci, M. Can, *Some results on q-analogue of the Lerch zeta function*, Adv. Stud. Contemp. Math. **12** (2006), 213–223.

[3] F. T. Howard, *Applications of a recurrence formula for the Bernoulli numbers*, J. Number Theory **52** (1995), 157–172.

[4] T. Kim, *A note on $q-$Volkenborn integration*, Proc. Jangjeon Math. Soc. **8** (2005), 13–17.

[5] T. Kim, *$q-$Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), 288–299.

[6] T. Kim, *A note on p-adic invariant integral in the rings of p-adic integers*, Advan. Stud. Contemp. Math. **13** (2006), 95–99.

[7] T. Kim, L.-C. Jang, H. K. Pak, *A note on q-Euler and Genocchi numbers*, Proc. Japan Acad. Ser. A Math. Sci. **77** (2001), 139–141.

[8] T. Kim, *A note on some formulas for the q-Euler numbers and polynomials*, Proc. Jangjeon Math. Soc. **9** (2006), 227–232.

[9] T. Kim, *On the twisted q-Euler numbers and polynomials associated with basic q-l-functions* , J. Math. Anal. Appl. (in press).

[10] T. Kim, *The modified q-Euler numbers and polynomials*, arXiv: math.NT/0702523 (2006).

[11] T. Kim, *An invariant p-adic q-integral on $\mathbb{Z}_p$*, Appl. Math. Lett.(2007), doi:10.1016/j.aml.

[12] H. M. Srivastava, T. Kim and Y. Simsek, *q-Bernoulli numbers and polynomials associated with multiple q-zeta functions and basic L-series*, Russ. J. Math. Phys. **12** (2005), 241–268.

Taekyun Kim
EECS, Kyungpook National University, Taegu 702-701, S. Korea
e-mail: tkim@knu.ac.kr; tkim64@hanmail.net